

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/373096862>

Conservation Agriculture: A Long-term Approach towards Sustainability

Article in *International Journal of Environment and Climate Change* · August 2023

DOI: 10.9734/IJECC/2023/v13i102625

CITATIONS

0

READS

22

6 authors, including:



Jyoti Sharma

Sher-e-Kashmir University of Agricultural Sciences and Technology Jammu

19 PUBLICATIONS 6 CITATIONS

[SEE PROFILE](#)



Monika Menia

Sher-e-Kashmir University of Agricultural Sciences and Technology Jammu

13 PUBLICATIONS 23 CITATIONS

[SEE PROFILE](#)



Deepak Kumar

Sher-e-Kashmir University of Agricultural Sciences and Technology Jammu

14 PUBLICATIONS 11 CITATIONS

[SEE PROFILE](#)



Ranjeet Singh Bochalya

MS Swaminathan School of Agriculture Shoolini University Solan

26 PUBLICATIONS 21 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



SKUAST-Jammu [View project](#)



Response of different sources of phosphorus on growth nodulation and yield of green gram (vigna radiata) [View project](#)



Conservation Agriculture: A Long-term Approach towards Sustainability

**Jyoti Sharma ^{a*}, Amit Mahajan ^{b++}, Monika Menia ^c,
Deepak Kumar ^d, Ranjeet Singh Bochalya ^e, Naveena ^a
and Shesh Narayan Kumawat ^a**

^a Division of Agronomy, Sher-e-Kashmir University of Agricultural Science and Technology of Jammu, Jammu and Kashmir, India.

^b Krishi Vigyan Kendra Samba, 184121, India.

^c Department of Agricultural Sciences, DAV University, Jalandhar, Punjab, India.

^d Department of Agriculture, Galgotias University, Greater Noida, Uttar Pradesh 203201, India.

^e M.S. Swaminathan School of Agriculture, Shoolini University of Bio-Technology and Management, Solan, H.P., 173229, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102625

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102918>

Review Article

Received: 14/05/2023

Accepted: 19/07/2023

Published: 12/08/2023

ABSTRACT

With the application of intensive agricultural techniques, conventional agriculture has been successful in meeting production goals but has also led to the depletion of natural resources. Sustainability in the management of the natural resource base is necessary for ongoing and expanded agricultural output to provide food security for future generations. Over a long period, the traditional tillage practice has led to the destruction of the natural resource base of the land. As a result of its inherent connection to the physical, chemical, and biological aspects of soil, it has led to a significant loss of soil and SOM, which is a crucial component of soil quality. Therefore, a suitable cropping system and land use must be implemented and adopted to ensure food security on a

⁺⁺ Programme Assistant (Farms);

^{*}Corresponding author: E-mail: sharma32jyoti@gmail.com, jyoti12121sharma@gmail.com;

sustainable basis. These measures should be based on principles to prevent land degradation, protect the natural resource base, and improve food and nutritional security through crop diversification and optimal rotation. The one sustainable cropping method that may reverse soil erosion, increase crop output, and improve the socioeconomic status of small landholder farmers is conservation agriculture (CA). Conservation agriculture (CA) is the integrated management of the available natural resources such as soil, water, flora, and fauna, with certain outside inputs to maximize the effectiveness of natural resource utilization. To fulfill the objective of sustainable crop production, an alternative approach known as conservation agriculture has emerged. It represents a significant advance in the direction of sustainable agriculture. This article reviews the emerging concerns due to the continuous adoption of conventional agriculture systems, contrasting features between conventional and conservation agriculture systems, and various principles and practices in conservation agriculture. It also highlights the benefits and limitations of CA and various challenges in the adoption of CA.

Keywords: *Conservation agriculture; crop diversification; production; natural resources; and sustainability.*

1. INTRODUCTION

By 2050, the population of the globe is expected to reach 9.1 billion people [1]. Based on the limited land resources, it is clear that urgent measures must be taken to assure enhanced food supply and food security. Agriculture, which is the sole source of human food, is the world's largest industry and major land use, accounting for 40% of all available land [2]. Since the global supply, economic growth, and access are not keeping up with the rising population in developing nations, food security has become more crucial on both the international and domestic fronts. Through industrialization and the expansion of agricultural production into marginal lands, rapid population increase and economic development gravely damaged the ecosystem. Growing issues with resource degradation, such as groundwater depletion, water logging, salinization, soil erosion, biodiversity loss, and invasive species, further compound the problem of food security [3]. The task of supplying food demand is made even more difficult by the declining per capita availability of arable land and the delayed pace of climate change adaptation. While climate change may have an impact on the entire planet, its effects are particularly severe in Asian regions due to the region's high reliance rate [4]. Therefore, the main challenges faced by most Asian countries are ensuring food security for a growing population and reducing poverty while maintaining agricultural systems in the face of depleting natural resources, adverse effects of climate variability, spiraling input costs, and unstable food prices. Along with these challenges, soil erosion, a reduction in soil organic matter, and salinization are the main warning signs that agricultural systems are not sustainable. These are mostly brought about by

(i) heavy tillage-induced soil organic matter reduction, soil structural degradation, water, and wind erosion, decreased rates of water infiltration, surface sealing and crusting, soil compaction, (ii) insufficient return of organic material, and (iii) monoculture. Low-input subsistence farming typically results in lower crop yields and deteriorating soil quality [5,6]. Excessive and improper tillage decreases soil productivity and soil organic carbon [7] and speeds up soil erosion and degradation [8,9,10]. Thus, intensification of agriculture is required without further depleting the natural resource base. Therefore, a paradigm change in farming practices is essential for future productivity advances while preserving natural resources (eliminating unsustainable components of conventional agriculture, such as plowing/tilling the soil, removing all organic material, and monoculture) [11]. The numerous crop management techniques referred to as "conservation agriculture" (CA) are widely used to increase crop output (Hernandez et al. 2018a; 2018b), conserve soil, and build robust systems to weather-induced pressures (Hernandez et al. 2018c), especially those brought on by irregular weather patterns (Hernandez et al. 2017) and climatic change [12]. CA is a significant agronomic practice that has steadily expanded globally to cover about 11% of the world's arable land (157.8 Mha) [13]. FAO (2010b), has defined CA as a concept for resource-saving agricultural productivity that aims to provide high and sustained levels of output while also preserving the environment. Dumanski et al. [14] stated that conservation agriculture (CA) is not "business as usual", as it does not aim to maximize yields while utilizing the resources of the soil and agro-ecosystem. Instead, CA focuses on increasing profitability and yields to strike a

balance between agricultural, economic, and environmental benefits. It argues that the social and economic gains from both production and environmental protection, including lower material and labor costs, outweigh those from production alone. By using fewer fossil fuels, pesticides, and other pollutants while preserving environmental services and integrity, farming communities can provide more hygienic living conditions for the larger population. The three main tenets of CA are systematic crop rotation, crop residues that permanently coat the soil, and minimal or no tillage [15]. The FAO guidelines state that "CA is an approach to managing agricultural ecosystems for enhanced and sustained productivity, improved returns, and food security while maintaining and improving the resource base and the environment." It is an agronomic practice that includes rotational planting of pulses and legumes together with reduced tillage (RT), no-tillage (NT), or minimum tillage with stable organic matter cover, and crop residue retention. It is stated that the aforementioned concepts apply to all CA systems [16]. However, elements that are specific in nature, such as the choice of farm implements, establishing techniques, rotation of crops with pulses or legumes, management of soil fertility, management of mulch and crop residue, etc., vary depending on the environment. The prospect of CA is that it may

be used in many agricultural ecological zones and beneficial in improving food security for a significant number of smallholder residents of underdeveloped nations [17]. Comparing Conservation Agriculture (CA) to conventional tillage (CT) systems, researchers found that CA can alter soil's physical, chemical, and biological soil quality metrics. In response, the increased sink for carbon storage within the soil helps to counterbalance the climate variability, which has an impact on ecosystem services and the sustainability of crop production systems (Yadav et al., 2017). He stated that conservation agriculture is a successful strategy for promoting an agricultural production system that includes a greater diversity of soil microorganisms, which are crucial for enhanced soil quality, crop yield, and numerous ecosystem services. For sustaining prospects, CA is the best management of natural resources like soil, water, vegetation, and biodiversity. CA have the ability to mitigate the consequences of climate change by increasing crop output and benefits while coordinating agricultural, financial, and ecological benefits [18].

Conventional and Conservation Agriculture:

To implement conservation agriculture systems, traditional agriculture must completely change the management of crops, land, water, nutrients, weeds, and farm machinery.

Table 1. Contrasting features between conventional and conservation agriculture

Conventional agriculture	Conservation agriculture
Cultivation using science and technology to dominate nature	Least interference with natural processes
Excessive mechanical tillage and soil erosion	No-till or significantly reduced tillage (biological tillage)
Monocultures or crop rotation in tillage framework	Crop rotation or intercropping is a permanent feature
Residue burning or removal (bare surface)	Surface retention of residues (permanently covered)
The oxidation of organic matter caused by traditional tillage's exposure to air and sunlight results in low soil carbon content, which has an impact on soil structure.	Accumulation of the organic matter in soil provides better soil aggregation.
Low infiltration rate of water	High infiltration rate of water
High wind and soil erosion	Low wind and soil erosion
Kills established weeds but also encourages more weed seeds to germinate	Weeds are a problem in the beginning but decrease over time
Use of ex-situ FYM/composts	Use of in-situ organics/composts; Brown manuring/cover crops (surface retention)
Free-wheeling of farm equipment and increased soil compaction.	Controlled traffic, tramline compaction, and reduced soil compaction in crop areas.

[19]

2. PRINCIPLES OF CONSERVATION AGRICULTURE

1. **Permanent or semi-permanent soil cover with organic materials:** A permanent soil cover is essential for three reasons: it protects the soil from the harmful consequences of exposure to rain and sunlight, it continuously supplies "food" to the micro- and micro-organisms in the soil, and it modifies the microclimate in the soil to promote the growth and development of soil organisms, including plant roots. Therefore, it enhances soil biological activity, biodiversity, soil aggregation, and carbon sequestration [20]. Kumar and Goh (2000) studied the impact of crop residues and management techniques on soil quality, soil nitrogen dynamics and recovery, and crop output and reported that agricultural residues from cultivated crops have a major impact on crop productivity due to their effects on the physical, chemical, and biological activities of the soil as well as the quality of the water and soil. When a previous crop is left anchored or loose after harvest or when a cover crop (legume or non-legume) is established and killed or clipped to provide mulch, crop residue is produced. Composts and manures can also be used as external mulch, but their use may be limited to higher-value crops like vegetables due to the cost of transporting such large materials to the field. The force of raindrops striking bare soil causes runoff and soil erosion by destroying soil aggregates, blocking soil pores, and drastically reducing water penetration. Mulch deflects this energy, preventing soil aggregate breakdown on the surface while also improving water infiltration and reducing soil erosion [21]. In comparison to tilled soils, no-tillage with mulch improves water infiltration, decreases runoff, and boosts yield [22]. Roldan et al. [23] demonstrated that after 5 years of NT maize in Mexico, soil wet aggregate stability had increased over conventional tillage, as had soil enzymes, soil organic carbon (SOC), and microbial biomass (MBM). Madari et al. [24] demonstrated that NT with residue cover had higher aggregate stability, higher aggregate size values, and total organic carbon in soil aggregates than TT in Brazil. Through competition and depriving weed seeds of the light they frequently need for germination, a cover crop and the subsequent mulch or leftover crop residue assist in reducing weed

infestation. Additionally, there is proof that cereal residues have allelopathic characteristics that prevent the germination of surface-level weed seeds. (Jung et al., 2004). Vagen et al. (2005), reported that the creation of natural or modified fallow systems (agroforestry) with attainable C accumulation rates of 0.1 to 5.3 Mg C ha⁻¹ yr⁻¹ has the greatest potential for raising SOC. According to them, adding crop wastes or manure to cropland in addition to NT can produce C accumulation rates that are attainable, up to 0.36 Mg C ha⁻¹ yr⁻¹. Lal (2005) predicted that increasing SOC by 1 Mg ha⁻¹ yr⁻¹ can enhance food grain production by 32 Mg yr⁻¹ in impoverished nations as SOC is a crucial measure of soil quality. Through their roots, cover crops aid in promoting biological soil tillage; the top mulch supplies food, nutrients, and energy to underground worms, arthropods, and microorganisms that also biologically till soils. Compaction under zero-tillage systems can also be reduced by using biological agents (earthworms, etc.) and deep-rooted cover crops. According to recent studies, more soil fauna is found in no-tillage, residue-retained management regimes compared to tillage plots. The networks of soil pores, particularly those of mycorrhizal hyphae, which are crucial for the availability of phosphorus in some soils, are damaged and disrupted by tillage [25]. Thus, zero-tillage produces healthier soil with a greater mix of bacteria and other species. Ground cover encourages an increase in biodiversity both below and above ground; ground cover and mulch have been shown to enhance the number of beneficial insects, which in turn helps control insect pests [26]. Crop health, yield, and soil quality are all impacted by interactions between rhizobacteria and root systems. Exudates released by plants stimulate and maintain certain rhizobacterial populations that improve nitrogen fixation, bio-control of plant pathogens, disease resistance, and nutrient cycling.

2. **Minimum Mechanical Soil Disturbance or No-Tillage (NT) or Reduced Tillage or Minimum Tillage:** The biological activity of the soil creates very solid soil aggregates and holes of different sizes that enable the infiltration of air and water. The term "biological tillage" might be used to describe this process, which is incompatible with mechanical tillage. The biological soil structuring mechanisms will cease to exist with mechanical soil disturbance. Minimal

soil disturbance promotes/maintains ideal proportions of respiration gases in the rooting zone, moderate organic matter oxidation, porosity for water transport, retention, and release, and restricts the exposure of weed seeds to new light and their germination [27]. Minimal tillage refers to either direct sowing or dissemination of seed with the least amount of soil disturbance by opening the slot with the use of a khurpi or other equipment. Regardless of what is lower, the disturbed area shall be no wider than 15 cm or no larger than 25% of the harvested area [28]. Numerous advantages of minimal soil disturbance were stated in the section on permanent soil cover above, and it is crucial to combine these two techniques for the greatest outcomes. When used for plowing, tractors generate greenhouse gases (mostly CO₂) that increase expenses while also consuming a significant amount of fossil fuels (Grace et al., 2003). Animal-based tillage systems are particularly costly because farmers must care for and feed a pair of animals for a whole year. Animals also release methane, a greenhouse gas that is 21 times more powerful than carbon dioxide in causing global warming (Grace et al., 2003). These expenses and emissions are decreased with zero-tillage. According to farmer surveys conducted in Pakistan and India, zero-tilling wheat after rice lowers production costs by US\$60 per hectare by using less fuel (60-80 l ha⁻¹) and manpower [29]. Since farmers utilize 2 to 12 passes of a plow to create a good seedbed, the turnaround time in this rice-wheat system from rice to wheat ranges from 2 to 45 days (Hobbs & Gupta 2003). With zero-till wheat, this period is only 1 day long. Due to increased oxidation brought on by tillage and modern agricultural practices, soil organic matter decreases with time, resulting in soil degradation and a loss of soil biological fertility and resilience [30]. On the other hand, zero-tillage has been demonstrated to cause an accumulation of organic carbon in the surface layers when paired with permanent soil cover (Lal, 2005). No-tillage is a potential method for maintaining or even increasing soil C and N stocks since it reduces SOM losses (Bayer et al., 2000). Leake [31] discusses the impact of tillage on different types of soil diseases using numerous instances. In his conclusion, he acknowledged that healthy soil with a high microbial diversity does play

a role by being hostile to soil pathogens and stated that the impact of tillage on diseases is uncertain. To help shift the advantage from the disease to the crop, he also recommended that NT farmers alter management to control infections by sowing date, rotation, and resistant cultivars. Tillage increases the amount of wear and tear on equipment and increases tractor maintenance costs relative to zero-tillage systems, which is another economic factor to take into account.

3. **Crop Rotation with Pulses or Legumes:**

Crop rotation is essential not just to provide a varied "diet" for soil microorganisms, but also to explore through various soil levels for nutrients that have leached to deeper layers and can be "recycled" by crops in rotation. A diversified crop rotation also results in diverse soil flora and fauna. Legumes are used in crop sequences and rotations because they disrupt the life cycles of pest species, fix nitrogen biologically, reduce off-site pollution, and increase biodiversity [27]. Rotation should comprise a minimum of three different crops. Crop rotation is a method of agricultural management that has a long history. In zero-till systems, a larger network of root channels and macro-pores is encouraged by the rotation of several crops with various rooting patterns and low soil disturbance. This aids in the penetration of water into deeper levels. The risk of pests and disease outbreaks caused by pathogenic organisms is decreased as a result of rotations because the biological diversity helps keep pathogenic organisms in check [31]. The life cycle of many weeds is disrupted by rotating crops, which also results in a decline in the overall weed population. Due to these advantages, crops cultivated in rotation often produce 10% more than crops grown in monoculture [32]. Additionally, a variety of crops grown in rotation produce a diversity of soil flora and fauna because the roots excrete various organic substances that draw various bacteria and fungi, which are crucial for the conversion of these substances into nutrients for plants [33]. It is crucial for soil health because it lessens the allelopathic effect of crops and improves soil fertility and quality while also increasing crop output. Monocropping or growing the same crop in the same location for an extended period hurts the health of the soil, making it sick. Diversified crops are cultivated, which

improve soil porosity, moisture retention, root penetration, and nutrient recycling [34]. It results in better N/P/K balance from both organic and mineral sources, as well as better biological N₂ fixation through specific plant-soil biota symbionts.

3. BENEFITS OF CONSERVATION AGRICULTURE

1. **Soil Aggregation, Aggregate Stability, and Structure:** The sensitivity of soil to alter under natural or manmade activity is the subject of soil aggregate stability. Because soil erodibility is closely correlated with aggregate stability, it is crucial for soil conservation owing to water erosion. Between aggregate stability in water, aggregate size, and total organic carbon content, there is a medium-to-high aggregate connection [35]. Conventional tillage encourages soil organic matter loss, which causes soil aggregate disintegration and erosion [35]. CA methods that leave more agricultural residue on the soil's surface typically allow for improvements in aggregate stability and soil aggregation. Additionally, it shields surface aggregates from splash or raindrop degradation. According to studies conducted in California, no-tillage with stubbles that retained treatment exhibited greater water-stable aggregation. No-tillage CA that retains crop residue is beneficial for soil aggregation and aggregate stability [36]. According to studies on soil structure, persistent no-tillage management in semi-arid Morocco promoted the growth and

persistence of a soil surface horizon rich in stable aggregates [37]. Under no-tillage as opposed to traditional tillage, Mrabet [38] found greater mean weight diameter (MWD) and aggregation index (AI) at the surface (0–7 cm) of a self-mulching expanding clay soil. To improve infiltration, plant-available water, and aggregate stability, CA needs a suitable amount of soil cover (Palm et al., 2013). Additionally, it raises the fraction of soil micropores, boosts water-holding capacity, and lowers evaporation.

2. **Soil and moisture conservation:** From the perspective of sustainability and off-site environmental harm, runoff, and soil loss are key concerns of sloping agriculture land in Himalayan regions, particularly those with unstable aggregates in the surface horizon. As such, this has been and continues to be an extremely important research field. CA with agricultural residue mulch can offer soil cover to lessen the impact of rain and provide barriers against runoff; this will aid in increasing moisture uptake and reducing soil separation (Franzluebbers 2002). According to research on the western Loess Plateau of China, a wheat-pea rotation system decreased soil loss from erosion by about 62% while reducing runoff and runoff intensity with no-tillage and stubble retention (Zhao et al. 2007). In summary, CA could lessen soil detachment and improve water infiltration, which implies a reduction in water flow; as a result, soil erosion would be decreased (Table 2). The fact that conventional tillage (CT) results in more




Fig. 1. Principles of conservation agriculture




Fig. 2. a) Plants growing in good soil structure b) Plants growing in compacted soil shows three types of compaction: surface crusting, plow layer/surface compaction, and subsoil compaction

Table 2. Different tillage systems, runoff, and infiltration

Tillage system	Five years means as % of rainfall	
	Seasonal runoff	Seasonal infiltration
Conventional tillage	20	80
Conservation tillage	1	99

[42]

physical disruption and less production of aggregate stabilizing elements in conventional farming systems leads to higher soil deterioration [39]. Additionally, adding crop residue through plowing, removing them from the field for use as cow feed, or burning them exposes the soil to the effects of rain, wind, and solar heating, speeding up the pace of soil degradation. Lower soil erosion potential in CA practices than in traditionally tilled areas is caused by higher aggregate stability in CA [40]. In contrast to CT, CA leaves more plant debris on the surface, protecting the soil from the damaging impacts of rain, strong winds, and solar heating. Due to less runoff in fields under CA circumstances, soil erosion is further declined [41].

3. **Enhance Soil Quality:** The ability of a particular type of soil to function, within naturally controlled ecosystem boundaries, in a way that supports plant and animal productivity, maintains or improves water and air quality, and supports human health

and habitation is known as soil quality [43]. Verhulst et al. [44] stated that In terms of agricultural output, excellent soil quality, "equates to the ability of the soil to maintain a high productivity without significant soil or environmental degradation." The physical, chemical, and biological aspects of the soil are used to evaluate its quality. Good-grade soil can be referred to as "healthy soil" in terms of biological soil quality (2). A stable system with high levels of biological diversity and activity, internal nutrient cycling, and resistance to disturbance is referred to as healthy soil [45]. When CA is used for a sufficient amount of time, the quality of the soil is significantly improved, especially in the surface layers [44]. When evaluating the sustainability of agricultural production systems, soil structure is an important component to consider [44]. It is frequently described as the degree of stability of aggregates [46]. In comparison to CT, ZT with residue retention increases both the dry and the wet aggregate size distribution [40].

While CA reduces soil compaction due to reduced tillage operation and growth of the deep-rooted cover crops or legumes in rotation, which break the compact layers in the sub-surface, CT, for example, can cause compactness in soil subsurface layers leading to restricted root growth, water logging, and poor aeration [30]. Recent studies showed that CA improves aeration and water retention by reducing bulk density, especially in surface layers [47]. In soils under CA, residue retention leads to increased microbial biomass and an abundance of earthworms and macroarthropods, such as termites and ants, which benefits soil fertility. Due to extensive nitrification, NO₃-leaching, and H₃O⁺ excretion by legume roots, the addition of legumes to crop rotations in CA may lower the pH of alkaline soils [48]. Thus, it can be said that soils under CA are generally layered physically, chemically, and biologically, with the surface layers having better soil quality.

4. **Enhance Nutrient Use Efficiency:** The distribution, recycling, and transformation of nutrients in soils are significantly influenced by conservation tillage, crop residue management, and crop rotation with pulses or legumes [44]. CA can affect the availability of soil nitrogen because of its effects on soil organic carbon and nitrogen mineralization [39,49]. Under CA, soil fertility often increases over time and requires fewer fertilizer amendments to produce ideal yields over time. Additionally, NT has been shown to increase the availability of potassium and phosphorus. Mrabet et al. [50] found that No-tillage soil has more phosphate and potassium near the soil surface than tilled soil. According to Rahman et al. [51], the surface soil under NT had much greater exchangeable Ca, Mg, and K levels than the soil that had been plowed. The biochemical decomposition of organic crop residue at the soil surface, which is significant for feeding the soil microbes, improves soil nutrient supply and cycling. Reduced runoff and the use of suitable deep-rooting cover crops help CA fields retain more nutrients [52]. Crop residues release nutrients gradually, preventing leaching and/or denitrification-related nutrient losses. Additionally, immobilizing mineral N through residue retention may help stop possible losses brought on by NO₃-N leaching [53]. The short-term immobilization of mineral nutrients

by microorganisms may result in reduced fertilizer use efficiency. Long-term nutrient availability, however, rises as a result of microbial activity and nutrient recycling [54].

5. **Enhance crop yields:** The initial soil fertility state [55], climate [56], the amount of rainfall experienced during the season [57], the management practices, and the type and quantity of crop residues kept, among other factors, all have an impact on the short-term effects of CA on crop production in comparison to CT [58,59]. As a result, the short-term impacts of CA on agricultural yield could be favorable, neutral, or adverse. However, over time, CA has reportedly been shown to increase crop yields because of its additional advantages, such as the reduction of soil erosion, higher soil quality, better moisture regimes, timely field activities (mostly sowing), and the advantages of crop rotation [60,61]. Mulching and rotational planting of legumes result in better soil physical, chemical, and biological qualities that, over time, minimize soil degradation and produce higher and more consistent yields in CA fields [62]. Due to enhanced RWUE from greater infiltration, less evaporation loss, higher soil water holding capacity, and timely crop planting in rainfed circumstances in dry areas where soil moisture is the main limiting factor, CA aids to boost agricultural yields. With the right fertilizer management, conservation agriculture may help boost crop output, enhance soil health, and generate revenue. Bell et al. [63] discovered that the longer-term practice of conservation agriculture enhanced rice grain output (by up to 12%). Khorami et al. [64] found that conservation agriculture, which includes reduced tillage, enhanced agronomy, and improved varieties, has produced beneficial results, including an increase in wheat yield and maize biomass. According to Baumhardt et al. [65], straw mulching increased the water flow of both rainfed and irrigated crops. Similarly, Bashour et al. [66] found that using conservation agriculture increased wheat crop yield by 27%. In southern Africa, conservation agriculture produced higher yields of maize (*Zea mays* L.) than conventional tillage, as reported by Thierfelder et al., [67]. Karki et al. (2014a) documented that conservation tillage with residue has a lower benefit-cost ratio (1.7) than conventional tillage without residue (2.5). Afzalnia et al. [68]

also observed increased wheat and maize (*Zea mays* L.) yield and biomass as a result of conservation agriculture. Mosquera et al. [69], reported that the adoption of a conservation agriculture system as opposed to conventional methods boosted crop yield and net (of the cost of production) benefits of the system by up to 25 and 24%, respectively. In potato-based systems, enhanced CA practices over 7 years revealed that soil carbon concentration in the entire profile was 29% higher under conservation tillage than under conventional tillage sites, and the carbon content was greater by 33% [70].

Table 3. Percent increase in grain yield in conservation agriculture

Crop	Increase in grain yield	References
Rice	12%	Bell et al. [63]
Wheat	27%	Gathala et al. [71]
Potato	24%	Mosquera et al. [69]
Lentil	27.7%	Bashour et al. [66]
Soyabean	20%	Thiagalingam et al. [72]
Bean	7%	Liben et al. [73]

6. **Enhance soil biodiversity:** When it comes to the activities of soil fauna and flora and the improvement of biological characteristics, conservation tillage is preferred above conventional tillage [74]. Cookson et al. [75] found that as tillage disturbance increased, bacterial biomass increased and fungal biomass decreased. Additionally, they discovered that the microbial community's composition and mode of substrate utilization had changed, with unique substrate usage in no-tillage soil. It offers a genuinely sustainable production system, preserving natural resources while also enhancing them [76]. It also increases the variety of soil biota, fauna, and flora (including wildlife), as well as the microbial count in agricultural production systems, all without compromising yields at high production levels. The conservation agriculture systems raise microbial biomass by 83%, lower the metabolic quotient by 32%, and enhance the MBC: total carbon ratio by 23%. CA increases agricultural output by disrupting insect and disease cycles, reducing weed growth, and enhancing nutrient cycling, soil fertility, and biodiversity of soil biota (Palm et

al., 2013). By providing habitats and food for birds, animals, reptiles, and insects among others, CA has been shown to promote above-ground biodiversity also (FAO, 2011).

7. **Mitigate the effects of climate change:** Conventional agriculture often emits more greenhouse gases (GHGs), which has a higher impact on climate change. Because of soil tilling, crop residue mixing, and biomass burning, conventional agriculture emits GHGs [77]. CA may reduce climate change by carbon sequestration, decreased CO₂ and N₂O emissions, and possibly reduced methane (CH₄) emissions. Due to SOM's slower breakdown and oxidation, less soil disturbance may increase carbon storage in CA crops [78]. By keeping crop residues on the soil's surface rather than burning them, like in CA, CO₂ emissions are decreased. Direct planting and avoiding tillage processes allow CA to save a significant amount of fuel, which lowers CO₂ emissions [29]. Long-term N₂O emission reduction in CA fields may be possible as a result of a decreased requirement for nitrogenous fertilizers brought on by better soil fertility. Conservation agriculture will boost the resilience of crops to adapt to local climate change through improved soil quality and nutrient cycling [77]. The main ways that CA aids in adaptation to climate change are through improved soil moisture status, reducing extreme soil temperatures [79,80,81], timely agricultural activities, and improved crop health in CA fields, thus minimizing the impacts of climate change [82,83].

4. LIMITATIONS OF CONSERVATION AGRICULTURE

- The initial high cost of specialized sowing and/or planting equipment and the need for technical know-how for improved management are two short-term drawbacks of conservation agriculture [84]
- Initial years of adoption result in lower yields, higher input costs, more labor-intensive weeding, competition between mulching and animal feed for crop residue, and promotion of the practice as a universal method without taking socioeconomic or agroecological factors into account [85].
- The absence of understanding about locally adapted cover crops that yield quality biomass under the existing conditions is a particularly significant knowledge gap.

- The adaptability and ingenuity of the farmer, extension agents, and researchers in a particular area are crucial factors that determine whether CA is successful or unsuccessful.

4.1 Challenges in Conservation Agriculture

To address the many, adaptable, and context-specific needs of technologies and their management, conservation agriculture, an emerging paradigm for growing crops, will need an innovative system viewpoint. Research and development (R&D) in conservation agriculture will therefore require several novel characteristics to address the issue. These include:

- (a) Being aware of the system:** Conservation agricultural systems are far more complicated than traditional systems. Site-specific information is the greatest obstacle to the CA system's widespread adoption [86,87]. Understanding fundamental processes and how components interact to build a functioning system will be crucial to manage these systems effectively. For instance, crop residues that are kept on the surface operate as mulch, which lowers soil water losses owing to evaporation and upholds a stable soil temperature regime [88]. Crop residue can be a readily decomposable source of organic matter, but it can also host unwanted pest populations or otherwise change the ecology of the system. No-tillage techniques will affect the root system's spread and depth of encroachment. Therefore, it is necessary to acknowledge conservation agriculture as a system and create management plans.
- (b) Burning crop residues:** Farmers prefer to seed the crop in time by burning the residue to sow the following crop promptly and without the need for machinery for sowing under CA systems. This is now a usual aspect of the rice-wheat system in north India. This causes environmental and health issues [89].
- (c) Developing a system and a farming system perspective:** A system perspective is developed by collaborating with farmers. Therefore, creating and promoting new technologies will require a core group of scientists, farmers, extension agents, and other stakeholders operating in partnership mode. The system is to determine research

goals and allocate resources within a framework in this case, which is rather different from typical agricultural R&D. Relationship-building and looking for connections with partners that operate in complementary disciplines are given less consideration [90].

- (d) Site specificity:** Adapting techniques for conservation agricultural systems will be very site-specific, but learning from different locations will be a valuable approach to comprehending why particular technology or practices are useful in some conditions but not in others. Building a foundation of knowledge for sustainable resource management will be accelerated by this learning approach.
- (e) Skilled and scientific staff:** Expert and scientific manpower is required for managing CA systems, and researchers' capacity to approach issues from a systems viewpoint and collaborate closely with farmers and other stakeholders must be improved. We urgently need to improve the mechanisms for exchanging knowledge and information [91-103].

5. CONCLUSION

Different from the usual framework for agricultural research and development, which was primarily focused on meeting specific food grain production targets in India, conservation agriculture offers a fresh approach. Given the pervasive issues of resource degradation that came along with previous tactics to boost production with little regard for resource integrity, a paradigm shift has become necessary. To achieve continuous productivity growth, it is now essential to integrate issues related to productivity, resource conservation, soil quality, and the environment. The knowledge base required for creating and promoting CA systems will be extremely difficult to meet. For this, scientists' ability to approach issues from a systems viewpoint, collaborate closely with farmers and other stakeholders, and promote knowledge and information-sharing must all be significantly increased. By lowering cultivation costs, and increasing resource use efficiency, competitiveness, and sustainability in agriculture, conservation agriculture provides a chance to halt and reverse the downward spiral of resource degradation. However, there is a need to support research in India to determine the best crop combinations, management techniques, weed management, insect pest, and disease control

tactics, as well as to create CA machinery that is appropriate for different agro-climatic areas. The new mission must be "Conserving resources - enhancing productivity."

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAO. Global agriculture towards 2050. Rome: Food and Agriculture Organization; 2009a.
2. Sah D, Devakumar AS. The carbon footprint of agricultural crop cultivation in India. *Carbon Manag.* 2018;9(3): 213-25.
DOI: 10.1080/17583004.2018.1457908
3. Oliver MA, Gregory PJ. Soil, food security and human health: A review. *Eur J Soil Sci.* 2015;66(2):257-76.
DOI: 10.1111/ejss.12216
4. Kumar MD. Food security and sustainable agriculture in India: The water management challenge. Vol. 60. Colombo, Sri Lanka: International Water Management Institute; 2003.
5. Orlando Olivares B, Lobo D, Rey JC, Vega A, Rueda MA. Relationships between the Visual Evaluation of Soil Structure (VESS) and soil properties in agriculture: A meta-analysis. *Sci Agropecu.* 2023;14(1):67-78.
DOI: 10.17268/sci.agropecu.2023.007
6. Montenegro-Gracia EJ, Pitti-Rodríguez JE, Olivares-Campos BO. Identification of the main subsistence crops of the Teribe: a case study based on multivariate techniques. *idea.* 2021;39(3):83-94.
DOI: 10.4067/S0718-34292021000300083
7. Lal R. Soil carbon sequestration to mitigate climate change. *Geoderma.* 2004;123(1-2):1-22.
DOI: 10.1016/j.geoderma.2004.01.032
8. Cerdà A, Flanagan DC, le Bissonnais Y, Boardman J. Soil erosion and agriculture. *Soil Till Res.* 2009;106(1):107-8.
DOI: 10.1016/j.still.2009.10.006
9. Olivares B, Verbist K, Lobo D, Vargas R, Silva O. Evaluation of the USLE model to estimate water erosion in an alfisol. *J Soil Sci Plant Nutr.* 2011;11(2):72-85.
DOI: 10.4067/S0718-95162011000200007
10. Olivares B, Lobo D, Verbist K. Application USLE model on erosion plots under soil conservation practices and water in San Pedro de Melipilla, Chile. *Rev Cienc Ingenieria.* 2015;36(1):3-10.
11. Bhan S, Behera UK. Conservation agriculture in India—Problems, prospects, and policy issues. *Int Soil Water Conserv Res.* 2014;2(4):1-12.
DOI: 10.1016/S2095-6339(15)30053-8
12. Vilorio JA, Olivares BO, García P, Paredes-Trejo F, Rosales A. Mapping projected variations of temperature and precipitation due to climate change in Venezuela. *Hydrology.* 2023;10(4):96.
DOI: 10.3390/hydrology10040096
13. FAO. Food and Agriculture Organization of the United Nations, Conservation agriculture; 2016.
14. Dumanski J, Peiretti R, Benites JR, McGarry D, Pieri C. The paradigm of conservation agriculture. *Proceedings of the world association of soil and water conservation.* 2006;1:58-64.
15. Rusinamhodzi L. Crop rotations and residue management in conservation agriculture. In: Farooq M, Siddique KHM, editors. *Conservation agriculture.* Springer International Publishing. 2015;21-37.
16. Sayre K. Effects of tillage, crop residue retention and nitrogen management on the performance of bed-planted, furrow irrigated spring wheat in northwest Mexico. Paper presented at the Conference of the International Soil Tillage Research Organization. 2000;15. Fort Worth.
17. Derpsch R, Friedrich T. Global overview of conservation agriculture adoption. In: *Lead Papers 4th World Congress on Conservation Agriculture.* New Delhi: World Congress on Conservation agriculture; 2009.
18. FAO. The State of food and agriculture. Women in agriculture. Closing the gender gap for development. Rome: Food and Agriculture Organization of the United Nations. 2011a;2.
19. Choudhary M, Ghasal PC, Kumar S, Yadav RP, Singh S, Meena VS, et al. Conservation agriculture and climate change: An overview. *Conserv Agric Approach Combat Clim Change Indian Himalaya.* 2016;1-37.
20. Ghosh PK, Das A, Saha R, Kharkrang E, Tripathy AK, Munda GC, et al. Conservation agriculture towards achieving food security in north East India. *Curr Sci.* 2010;99(7):915-21.
21. Dormaar JF, Carefoot JM. Implications of crop residue management and

- conservation tillage on soil organic matter. *Can J Plant Sci.* 1996;76(4):627-34.
DOI: 10.4141/cjps96-112
22. Thierfelder C, Amézquita E, Stahr K. Effects of intensifying organic manuring and tillage practices on penetration resistance and infiltration rate. *Soil Till Res.* 2005;82(2):211-26.
DOI: 10.1016/j.still.2004.07.018
23. Roldán A, Caravaca F, Hernández MT, García C, Sánchez-Brito C, Velásquez M, et al. No-tillage, crop residue additions, and legume cover cropping effects on soil quality characteristics under maize in Patzcuaro watershed (Mexico). *Soil Till Res.* 2003;72(1):65-73.
DOI: 10.1016/S0167-1987(03)00051-5
24. Madari B, Machado PLOA, Torres E, de Andrade AG, Valencia LIO. No tillage and crop rotation effects on soil aggregation and organic carbon in a Rhodic Ferralsol from southern Brazil. *Soil Till Res.* 2005;80(1-2):185-200.
DOI: 10.1016/j.still.2004.03.006
25. McGonigle TP, Miller MH. Mycorrhizae, phosphorus absorption, and yield of maize in response to tillage. *Soil Sci Soc Am J.* 1996;60(6):1856-61.
DOI:10.2136/sssaj1996.03615995006000060034x
26. Jaipal S, Singh S, Yadav A, Malik RK, Hobbs PR. Species diversity and population density of macrofauna of rice-wheat cropping habitat in semi-arid subtropical northwest India in relation to modified tillage practices of wheat sowing. In: Malik RK, Balyan RS, Yadav A, Pahwa SK, editors. *Herbicide-resistance management and zero-tillage in the rice-wheat cropping system. proceedings international workshop, Hissar, India.* 2002;166-71.
27. Kassam AH, Friedrich T. Perspectives on nutrient management in conservation agriculture. Invited paper. New Delhi, India: IV World Congress on Conservation Agriculture; 2009.
28. FAO. CA adoption worldwide, FAO-CA website; 2014b.
29. Hobbs APR, Gupta R. Problems and challenges of no-till farming for the rice-wheat systems of the Indo-gangetic plains in south. *Sustain Agric Int Rice-Wheat Syst.* 2004;123-42.
30. Lal R, Mahboubi AA, Fausey NR. Long-term tillage and rotation effects on properties of a central Ohio soil. *Soil Sci Soc Am J.* 1994;58(2):517-22.
DOI:10.2136/sssaj1994.03615995005800020038x
31. Leake AR. Integrated pest management for conservation agriculture. *Conserv Agric Environ Farmers Experiences Innov Socio-economy Policy.* 2003;271-9.
32. Vanlauwe B, Wendt J, Giller KE, Corbeels M, Gerard B, Nolte C. A fourth principle is required to define conservation agriculture in sub-Saharan Africa: the appropriate use of fertilizer to enhance crop productivity. *Field Crops Res.* 2014;155:10-3.
DOI: 10.1016/j.fcr.2013.10.002
33. FAO. Food and Agriculture Organization of the United Nations, conservation agriculture; 2015.
34. Sharma J, Sharma BC, Bharti V, Kumar R, Sharma A, Jamwal S. Novel resource conservation technology for increasing the production and productivity of rice-wheat cropping system in Indo-Gangetic plains of India. *J Ecofriendly Agric.* 2022;17(2):235-42.
DOI: 10.5958/2582-2683.2022.00047.8
35. Olivares BO, Calero J, Rey JC, Lobo D, Landa BB, Gómez JA. Correlation of banana productivity levels and soil morphological properties using regularized optimal scaling regression. *CATENA.* 2022a;208:105718.
DOI: 10.1016/j.catena.2021.105718
36. Li LL, Huang GB, Zhang RZ, Bill B, Guangdi L, Kwong YC. Benefits of conservation agriculture on soil and water conservation and its progress in China. *Agric Sci China.* 2011;10(6):850-9.
DOI: 10.1016/S1671-2927(11)60071-0
37. Mrabet R, El Brahli A. Soil and crop productivity under contrasting tillage management systems in semiarid Morocco. III World Congress on Conservation Agriculture: "Linking Production, Livelihoods and Conservation". Nairobi:6; 2005.
38. Mrabet R. Stratification of soil aggregation and organic matter under conservation tillage systems in Africa. *Soil Till Res.* 2002;66(2):119-28.
DOI: 10.1016/S0167-1987(02)00020-X
39. Bradford JM, Peterson GA. Conservation tillage. *Handbook of soil science.* 2000;247-70.
40. Govaerts B, Sayre KD, Lichter K, Dendooven L, Deckers J. Influence of permanent raised bed planting and residue

- management on physical and chemical soil quality in rain fed maize/wheat systems. *Plant Soil*. 2007;291(1-2):39-54.
DOI: 10.1007/s11104-006-9172-6
41. Araya T, Cornelis WM, Nyssen J, Govaerts B, Getnet F, Bauer H, et al. Medium-term effects of conservation agriculture-based cropping systems for sustainable soil and water management and crop productivity in the Ethiopian Highlands. *Field Crops Res*. 2012;132:53-62.
DOI: 10.1016/j.fcr.2011.12.009
42. Marongwe LS, Nyagumbo I, Kwazira K, Kassam A, Friedrich T. Conservation agriculture and sustainable crop intensification: A Zimbabwe Case Study. 2012;17:156-63.
43. Karlen DL, Mausbach MJ, Doran JW, Cline RG, Harris RF, Schuman GE. Soil quality: A concept, definition, and framework for evaluation (a guest editorial). *Soil Sci Soc Am J*. 1997;61(1):4-10.
DOI:10.2136/sssaj1997.03615995006100010001x
44. Verhulst N, Govaerts B, Verachttert E, Castellanos-Navarrete A, Mezzalama M, Wall P, et al. Conservation agriculture, improving soil quality for sustainable production systems. *Adv Soil Sci Food Sec Soil Qual*. 2010;17:137-208.
45. Shaxson F, Kassam AH, Friedrich T, Boddey B, Adekunle A. Underpinning conservation agriculture's benefits: The roots of soil health and function. In: Main background document for the Workshop on Investing in Sustainable Crop Intensification: The Case for Improving Soil Health. 2008;22-4.
46. Bronick CJ, Lal R. Soil structure and management: A review. *Geoderma*. 2005;124(1-2):3-22.
DOI: 10.1016/j.geoderma.2004.03.005
47. Nurbekov A. Manual on conservation agriculture practices in Uzbekistan. Tashkent, Uzbekistan. 2008;40.
48. Qadir M, Oster JD, Schubert S, Noble AD, Sahrawat KL. Phytoremediation of sodic and saline-sodic soils. *Adv Agron*. 2007; 96:197-247.
DOI: 10.1016/S0065-2113(07)96006-X
49. Olivares BO, Araya-Alman M, Acevedo-Opazo C, Rey JC, Cañete-Salinas P, Kurina FG, et al. Relationship between soil properties and banana productivity in the two main cultivation areas in Venezuela. *J Soil Sci Plant Nutr*. 2020;20(4):2512-24.
DOI: 10.1007/s42729-020-00317-8
50. Mrabet R, Ibno-Namr K, Bessam F, Saber N. Soil chemical quality changes and implications for fertilizer management after 11 years of no-tillage wheat production systems in semiarid Morocco. *Land Degrad Dev*. 2001;12(6):505-17.
DOI: 10.1002/ldr.464
51. Rahman MH, Okubo A, Sugiyama S, Mayland HF. Physical, chemical and microbiological properties of an andisol as related to land use and tillage practice. *Soil Till Res*. 2008;101(1-2):10-9.
DOI: 10.1016/j.still.2008.05.006
52. FAO. Conservation agriculture case studies in Latin America and Africa [introduction]. FAO Soils Bulletin No. 78; 2001.
53. Thomas GA, Dalal RC, Standley J. No-till effects on organic matter, pH, cation exchange capacity and nutrient distribution in a luvisol in the semi-arid subtropics. *Soil Till Res*. 2007;94(2):295-304.
DOI: 10.1016/j.still.2006.08.005
54. Carpenter-Boggs L, Stahl PD, Lindstrom MJ, Schumacher TE. Soil microbial properties under permanent grass, conventional tillage, and no-till management in South Dakota. *Soil Till Res*. 2003;71(1):15-23.
DOI: 10.1016/S0167-1987(02)00158-7
55. Olivares BO. Determination of the potential influence of soil in the differentiation of productivity and in the classification of susceptible areas to banana wilt in Venezuela [doctoral dissertation], Universidad de Córdoba, UCOPress. Spain; 2022.
56. Parra R, Olivares B, Cortez A, Rodríguez MF. Patrones de homogeneidad pluviométrica en estaciones climáticas del estado Anzoátegui, Venezuela. *Rev Multiciencias*. 2012;12 (Extraordinario):11-7.
57. Olivares Campos BO, Paredes F, Rey JC, Lobo D, Galvis-Causil S. The relationship between the normalized difference vegetation index, rainfall, and potential evapotranspiration in a banana plantation of Venezuela. *Sains Tanah J Soil Sci Agroclimatology*. 2021;18(1):58-64.
DOI: 10.20961/stjssa.v18i1.50379
58. López-Beltrán M, Olivares B, Lobo-Luján D. Changes in land use and vegetation in the agrarian community Kashaama, Anzoátegui, Venezuela: 2001-2013. *Rev Geogr Am Cent*. 2019;2(63):269-91.
DOI: 10.15359/rgac.63-2.10

59. López M, Olivares B. Normalized Difference Vegetation Index (NDVI) applied to the agricultural indigenous territory of Kashaama, Venezuela. UNED Research [journal]. 2019;11(2):112-21. DOI: 10.22458/urj.v11i2.2299
60. Sharma J, Bharti V, Kumar D, Menia M, Bochalya RS. Resource conservation technology for enhancing the rice productivity in India: a review. The Pharma Innovation [journal], SP-10(9). 2021;163-7.
61. Hernandez R, Olivares B, Arias A, Molina JC, Pereira Y. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. Idesia. 2020;38(2):95-102. DOI: 10.4067/S0718-34292020000200095
62. Sisti CPJ, dos Santos HP, Kohhann R, Alves BJR, Urquiaga S, Boddey RM. Change in carbon and nitrogen stocks in soil under 13 years of conventional or zero tillage in southern Brazil. Soil Till Res. 2004;76(1):39-58. DOI: 10.1016/j.still.2003.08.007
63. Bell R, Haque M, Jahiruddin M, Rahman M, Begum M, Miah M, et al. Conservation agriculture for rice-based intensive cropping by smallholders in the eastern Gangetic plain. Agriculture. 2019;9(1): 1-17. DOI: 10.3390/agriculture9010005
64. Khorami SS, Kazemeini SA, Afzalnia S, Gathala MK. Changes in soil properties and productivity under different tillage practices and wheat genotypes: A short-term study in Iran. Sustainability. 2018;10(9):3273. DOI: 10.3390/su10093273
65. Baumhardt RL, Schwartz R, Howell T, Evett SR, Colaizzi P. Residue management effects on water use and yield of deficit irrigated corn. Agron J. 2013;105(4):1035-44. DOI: 10.2134/agronj2012.0362
66. Bashour I, Al-Ouda A, Kassam A, Bachour R, Jouni K, Hansmann B, et al. An overview of Conservation Agriculture in the dry Mediterranean environments with a special focus on Syria and Lebanon. AIMS Agric Food. 2016;1(1):67-84. DOI: 10.3934/agrfood.2016.1.67
67. Thierfelder C, Matemba-Mutasa R, Rusinamhodzi L. Yield response of maize (*Zea mays* L.) to conservation agriculture cropping system in Southern Africa. Soil Till Res. 2015;146:230-42. DOI: 10.1016/j.still.2014.10.015
68. Afzalnia S, Karami A, Alavimanesh SM. Comparing conservation and conventional tillage methods in the corn wheat rotation. In: Power and machinery. International Conference of Agricultural Engineering-CIGR-AgEng.: agriculture and engineering for a healthier life, Valencia, Spain; 2012.
69. Mosquera VHB, Delgado JA, Alwang JR, López LOE, Ayala YEC, Andrade JMD, et al. Conservation agriculture increases yields and economic returns of potato, forage, and grain systems of the Andes. Agron J. 2019;111(6):2747-53. DOI: 10.2134/agronj2019.04.0280
70. Quintero M, Comerford NB. Effects of conservation tillage on total and aggregated soil organic carbon in the Andes. Open J Soil Sci. 2013;03(8): 361-73. DOI: 10.4236/ojss.2013.38042
71. Gathala MK, Ladha JK, Kumar V, Saharawat YS, Kumar V, Sharma PK, et al. Tillage and crop establishment affects sustainability of South Asian rice-wheat system. Agron J. 2011;103(4):961-71. DOI: 10.2134/agronj2010.0394
72. Thiagalingam K, Gould N, Watson P. Effect of tillage on rainfed maize and soybean yield and the nitrogen fertilizer requirements for maize. Soil Till Res. 1991;19(1):47-54. DOI: 10.1016/0167-1987(91)90108-A
73. Liben FM, Hassen SJ, Weyesa BT, Wortmann CS, Kim HK, Kidane MS, et al. Conservation agriculture for maize and bean production in the central rift valley of Ethiopia. Agron J. 2017;109(6):2988-97. DOI: 10.2134/agronj2017.02.0072.
74. Busari MA, Kukal SS, Kaur A, Bhatt R, Dulazi AA. Conservation tillage impacts soil, crop and the environment. Int Soil Water Conserv Res. 2015;3(2):119-29. DOI: 10.1016/j.iswcr.2015.05.002
75. Cookson WR, Murphy DV, Roper MM. Characterizing the relationships between soil organic matter components and microbial function and composition along a tillage disturbance gradient. Soil Biol Biochem. 2008;40(3):763-77. DOI: 10.1016/j.soilbio.2007.10.011
76. Hernández R, Olivares B. Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. Trop Subtrop Agroecosystems. 2020; 23(2):1-12.
77. Hobbs PR, Govaerts B. How conservation agriculture can contribute to buffering

- climate change. *Clim Change Crop Prod.* 2010;177-99.
78. Jat RA, Wani SP, Sahrawat KL. Conservation agriculture in the semi-arid tropics. *Adv Agron.* 2012b;117:191-273. DOI: 10.1016/B978-0-12-394278-4.00004-0
79. Olivares BO, Zingaretti ML, Demey Zambrano JA, Demey JR. Typification of agricultural production systems and the perception of climate variability in Anzoátegui, Venezuela. *FAVE Agricultural Sciences Section.* 2016;15(2):39-50. DOI: 10.14409/fa.v15i2.6587
80. Cortez A, Olivares B, Parra R, Lobo D, Rodríguez MF, Rey JC. Description of meteorological drought events in localities of the central mountain range, Venezuela. *Science, engineering and applications.* 2018;22-44:I(1). DOI: 10.22206/cyap.2018.vlii.pp23-45
81. Casana S, Olivares B. Evolution and trend of surface temperature and wind speed (1994-2014) at the Parque Nacional Doñana, Spain. *Rev Fac Agron (LUZ).* 2020;37(1):1-25.
82. Olivares B. Tropical conditions of seasonal rain in the dry-land agriculture of Carabobo, Venezuela. *La Granja. J Life Sci.* 2018;27(1):86-102. DOI: 10.17163/lgr.n27.2018.07
83. Zingaretti ML, Olivares B. Analysis of the meteorological drought in four agricultural locations of Venezuela by the combination of multivariate methods. *UNED Research [journal].* 2018;10(1):181-92. DOI: 10.22458/urj.v10i1.2026
84. Baudron F, Andersson JA, Corbeels M, Giller KE. Failing to yield? Plows, conservation agriculture and the problem of agricultural intensification: an example from the Zambezi valley, Zimbabwe. *J Dev Stud.* 2012;48(3):393-412. DOI: 10.1080/00220388.2011.587509
85. Corbeels M, De Graaff J, Ndah TH, Penot E, Baudron F, Naudin K, et al. Understanding the impact and adoption of conservation agriculture in Africa: A multi-scale analysis. *Agric Ecosyst Environ.* 2014;187:155-70. DOI: 10.1016/j.agee.2013.10.011
86. Olivares B. Systematization of traditional knowledge and ancestral ethnicity kariña in Anzoátegui state, Venezuela. *Rev Investig.* 2014;82(38):89-102.
87. Camacho R, Olivares B, Avendaño N. Agri-food landscapes: an analysis of the livelihoods of indigenous Venezuelans. *Rev Investig.* 2018;42(93):130-53.
88. Gupta R, Jat ML. Conservation agriculture: Addressing emerging challenges of resource degradation and food security in South Asia. *Conserv Agric.* 2010;1-18.
89. Tripathi S, Singh RN, Sharma S. Emissions from crop/biomass residue burning risk to atmospheric quality. *Int Res J Earth Sci.* 2013;1(1):24-30.
90. Abrol IP, Sangar S. Sustaining Indian agriculture—conservation agriculture the way forward. *Curr Sci.* 2006;91(8):1020-5.
91. Dangour AD, Lock K, Hayter A, Aikenhead A, Allen E, Uauy R. Nutrition-related health effects of organic foods: A systematic review. *Am J Clin Nutr.* 2010;92(1):203-10. DOI: 10.3945/ajcn.2010.29269, PMID 20463045
92. Derpsch R. Keynote: Frontiers in conservation tillage and advances in conservation practice. In: Stott DE, htar M, RH, Steinhart GC, editors. *Sustaining the global farm. Selected papers from the 10th International Soil Conservation Organisation;* 1999;24-9.
93. FAO. Conservation agriculture and sustainable crop intensification in Lesotho. *Integrated crop management.* Rome, Italy. 2010;10.
94. Hernández R, Pereira Y, Molina JC, Coelho R, Olivares B, Rodríguez K. Calendario de siembra para las zonas agrícolas del estado Carabobo en la República Bolivariana de Venezuela. Sevilla, Spain, editorial universidad internacional de andalucía. 2017;247.
95. Hernández R, Olivares B, Arias A, Molina JC, Pereira Y. Agroclimatic zoning of corn cultivation for the sustainability of agricultural production in Carabobo, Venezuela. *Rev Univ Geogr.* 2018a;27(2):139-59.
96. Hernández R, Olivares B, Arias A, Molina JC, Pereira Y. Identificación de zonas agroclimáticas potenciales para producción de cebolla (*Allium cepa* L.) en Carabobo, Venezuela. *J Selva Andina Biosph.* 2018b;6(2):70-82.
97. Hernández R, Olivares B, Coelho R, Molina JC, Pereira Y. Spatial analysis of the water index: an advance in the adoption of sustainable decisions in the agricultural territories of Carabobo, Venezuela. *Rev Geogr Am Cent.* 2018c; 60(1):277-99. DOI: 10.15359/rgac.60-1.10

98. Karki TB, Gadal N, Shrestha J. Studies on the conservation agriculture based practices under maize (*Zea mays* L.) based system in the hills of Nepal. Int J Appl Sci Biotechnol. 2014;2(2):185-92.
DOI: 10.3126/ijasbt.v2i2.10353
99. Kumar K, Goh KM. Crop residues and management practices: Effects on soil quality, soil nitrogen dynamics, crop yield, and nitrogen recovery. Adv Agron. 1999; 68:197-319.
DOI: 10.1016/S0065-2113(08)60846-9
100. Lal R. Enhancing crop yields in the developing countries through restoration of the soil organic carbon pool in agricultural lands. Land Degrad Dev. 2006;17(2):197-209.
DOI: 10.1002/ldr.696
101. Olivares BO, Rey JC, Perichi G, Lobo D. Relationship of microbial activity with soil properties in banana plantations in Venezuela. Sustainability. 2022b;14(20): 13531.
DOI: 10.3390/su142013531
102. Palm C, Blanco-Canqui H, DeClerck F, Gatere L, Grace P. Conservation agriculture and ecosystem services: an overview. Agric Ecosyst Environ. 2014; 187:87-105.
DOI: 10.1016/j.agee.2013.10.010
103. Vågen T-G, Lal R, Singh BR. Soil carbon sequestration in sub-Saharan Africa: A review. Land Degrad Dev. 2005;16(1): 53-71.
DOI: 10.1002/ldr.644

© 2023 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/102918>